

**UNCLASSIFIED**

---

**AD 268 123**

*Reproduced  
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY  
ARLINGTON HALL STATION  
ARLINGTON 12, VIRGINIA**



---

**UNCLASSIFIED**

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

SECRET  
BY ASIA  
268123

TECHNICAL REPORT No 47

# IONIC ASSOCIATION AND CORRELATION BETWEEN DOUBLE LAYER STRUCTURE AND ELECTRODE KINETICS

by

PAUL DELAHAY AND AKIKO ARAMATA <sup>1</sup>

Prepared for Publication  
in the  
Journal of Physical Chemistry

62-1-5

**XEROX**

Louisiana State University  
Department of Chemistry, Baton Rouge, Louisiana

ASTIA  
RECEIVED  
DEC 19 1961  
RECEIVED  
TIPDE

October 1961

Reproduction in whole or in part permitted for any  
purpose of the United States Government

The possibility of ion pair formation solely in the double layer region at the electrode in charge transfer reactions was suggested in a previous paper<sup>2</sup> from this Laboratory. It was discussed in some detail by Frumkin<sup>3</sup> to account for certain features of current-potential curves for the reduction of anions. Ion pair

---

(1) Graduate student since September 1960.

(2) P. Delahay and G.C. Mattax, J. Am. Chem. Soc., **76**, 5314 (1954).

(3) (a) A.N. Frumkin, Trans. Faraday Soc., **55**, 156 (1959). (b) A.N. Frumkin, Transactions of the Symposium on Electrode Processes, E. Yeager, ed., Wiley, New York, N.Y., 1961, p.1.

---

formation outside the double layer was not considered. Formation of ion pairs in the bulk of the solution was discussed recently by Joshi and Parsons<sup>4</sup>, together with other interpretations, to account

---

(4) K.M. Joshi and R. Parsons, Electrochim. Acta, **4**, 129 (1961).

---

for the properties of the electrical double layer at the interface between mercury and mixtures of barium chloride and hydrochloric acid. The influence of ion pair formation in the bulk of solution in the correlation between double layer structure and electrode kinetics does not appear to have been considered thus far and is investigated here. Results are given for the reduction of iodate on mercury in alkaline solutions of varying composition. This electrode reaction is advantageous because it occurs in a range of potentials in which there is very little specific adsorption of the ions involved. Application is made of the corrected Tafel plot recently mentioned in a previous paper from this Laboratory.<sup>5</sup>

---

(5) K. Asada, P. Delahay and A.K. Sundaram, J. Am. Chem. Soc., **83**, 3396 (1961).

---

### EXPERIMENTAL

Experimental methods were quite conventional and have been reported previously<sup>5</sup>. Adsorbable impurities (organic) were removed by treatment with activated charcoal as noted before<sup>5</sup>. The solution of KOH was titrated. The  $K_4Fe(CN)_6$  solution was prepared from the dehydrated salt and was titrated for verification<sup>6</sup>.

---

(6) I.M. Kolthoff and R. Belcher, "Volumetric Analysis", Vol. III, Interscience Publishers, Inc., New York, N.Y., 1957, p.135.

---

A five-compartment cell was used with the following electrodes: hydrogen electrode in I; D.M.E. or capillary of Lippmann electrometer and platinum cylinder for differential capacity measurements in III; a high-resistance saturated calomel electrode inserted in IV; and a mercury pool in V. Junctions between compartments were: a bridge with fritted glass disks for I-II; a 1 mm.-bore capillary for II-III; and fritted glass disks for III-IV and IV-V. Solutions in the compartments were: the solution being studied in II, III and IV; this solution, but without potassium iodate, in I and the bridge; and KOH in V. All potentials were measured against the hydrogen electrode in I, the electrode in V being the anode of the electrolysis circuit. The potential of the hydrogen electrode was verified against the S.C.E.:  $E = -0.964$  V. for 5 mM  $KIO_3$  + 20 mM KOH;  $-0.960$  V, for the same solution with 0.4 M KCl;  $-0.963$  V. for the solution with

0.2 M  $K_2SO_4$  instead of KCl; and -0.966 V. for the solution with 0.1 M  $K_4Fe(CN)_6$  instead of KCl or  $K_2SO_4$ .

Currents did not exceed 10% of the diffusion current and were corrected for mass transfer polarization by use of the factor  $\left[1 - (1/i_d)\right]$ . This correction factor is sufficiently accurate for maximum currents during drop life at the foot of a polarographic wave<sup>7</sup>.

---

(7) J. Koutecky and J. Cizek, Coll. Czechoslov. Chem. Commun., 21, 836 (1956).

---

#### DESCRIPTION AND DISCUSSION OF RESULTS

Parallel linear Tafel plots were obtained in the current density range,  $I = 20$  to  $800 \mu A.cm^{-2}$ , for the reduction of potassium iodate in the electrolytes of Table I. These plots yielded  $\alpha n_a = 0.75$  ( $\alpha$ , transfer coefficient;  $n_a$  number of electrons in the activated step) in good agreement with the previously reported<sup>8</sup> value of 0.77. Corrected Tafel plots were prepared according to

$$\ln I + \frac{zF}{RT} \Delta\varphi = \ln I^0 + \frac{\alpha n_a F}{RT} E_e - \frac{\alpha n_a F}{RT} (E - \Delta\varphi) \quad (1)$$

---

(8) M. Breiter, M. Kleinerman and P. Delahay, J. Am. Chem. Soc., 80, 5111 (1958).

---

where  $I$  is the current density,  $I^0$  the exchange current density,  $E$  the electrode potential,  $E_e$  the equilibrium electrode potential,  $\Delta\varphi$  the difference of potential across the diffuse double layer

from the plane of closest approach to solution,  $z$  the ionic valence with sign of the reduced species (iodate), and  $F$ ,  $R$  and  $T$  have their usual significance<sup>9</sup>. Values of  $\Delta\varphi$  were calculated by a previously

(9) See ref.5 for a discussion.

described method<sup>5</sup> from the theory of the diffuse double layer, experimental differential capacities<sup>10</sup> and the point of zero charge.

(10) Capacities were measured with  $\text{KIO}_3$  present as the reaction is so irreversible that the faradaic impedance is very large in comparison with the double layer impedance.

Parallel linear plots were obtained which yielded  $\alpha n_a = 0.89$  and the intercepts in Table I. The lines without and with 0.4 M  $\text{KCl}$  are essentially identical within the experimental error of 3-5 mV but the double layer correction is less satisfactory for 0.2 M  $\text{K}_2\text{SO}_4$  and, especially, 0.1 M  $\text{K}_4\text{Fe}(\text{CN})_6$ . Disagreement for  $\text{K}_2\text{SO}_4$  was previously noted<sup>8</sup> (without the use of corrected Tafel plots).

Several causes can be invoked for the disagreement between theory and experiment: variation of activity coefficient, ion pair formation in the double layer, influence of ion size, etc. (cf. ref.4). Further, the calculation of  $\Delta\varphi$  is based on the Gouy-Chapman theory, i.e. on a very simplified model not unlike that of the Debye-Hückel model for electrolytes. It is surprising, in this respect, that the double layer correction holds so well for concentrations as high as 0.4 M  $\text{KCl}$ <sup>11</sup>. Ionic association in the bulk of the solution, which

(11) P. Delahay, "Transaction of the Symposium on Electrode Processes", E. Yeager editor, John Wiley & Sons, New York, 1961, p.138.

is not negligible for  $K_2SO_4$  and  $K_4Fe(CN)_6$ , is also to be reckoned with. One has  $\log K = 0.96$ <sup>12a</sup>, for formation of  $K SO_4^-$ , and  $\log K = 2.3$ <sup>12b</sup> for formation of  $K^+Fe(CN)_6^{3-}$ . Ionic association seems

---

(12) (a) J. Bjerrum, G. Schwarzenbach and L.G. Sillen, "Stability constants, Part II", The Chemical Society, London, 1958, p.80.

(b) Ibid., p.32. (c) Ibid., pp.2 and 93.

---

negligible for KOH and KCl<sup>12c</sup>. The corresponding error on ionic concentrations, if ionic association is not considered, is significant, especially for  $K^+$  ions which are non-specifically adsorbed in the diffuse double layer at the potentials considered here. The corrected Tafel plots of Fig.1 were obtained after allowance was made for ionic association. The discrepancy for  $K_2SO_4$  noted above disappears and, in fact, there seems to be a slight overcorrection. Results for KOH alone, KOH + KCl, and KOH +  $K_2SO_4$  are nearly the same within experimental error. The discrepancy for  $K_4Fe(CN)_6$  is decreased but not entirely eliminated. This discrepancy, as given by Fig.1, is greater than it actually is because formation of the triple ion  $(K_2^+)Fe(CN)_6^{2-}$  was not considered. There is evidence for the formation of this ion<sup>13</sup> but no quantitative datum is available. Further, other

---

(13) S.R. Cohen, Thesis, Cornell University, Ithaca, 1956, p.34.

---

effects, and particularly the influence of ionic size could be invoked to explain the remaining small departure from theory. This matter will not be taken here, though it is being studied at the present in this Laboratory.



Ionic association of the reactant,  $\text{KIO}_3$  ( $K = -0.25$  for formation<sup>14</sup> of  $\text{K}^+\text{IO}_3^-$ ) should also be considered since the faradaic current is the sum of the currents for the reduction of  $\text{IO}_3^-$  and  $\text{K}^+\text{IO}_3^-$ . These

---

(14) Cf. ref.12, p.124.

---

two parallel electrode processes undoubtedly have different exchange densities. Further, kinetics of dissociation of  $\text{K}^+\text{IO}_3^-$  complicates matter. Double layer corrections for this type of process were discussed semi-quantitatively<sup>8, 15</sup>, and theoretical analyses have been worked out<sup>16, 17</sup>. This effect is probably minor here because of the

---

(15) L. Gierst, ref.11, pp.109-138.

(16) H. Matsuda, J. Phys. Chem., **64**, 336 (1960).

(17) H. Hurwitz, Z. Elektrochem., **65**, 178 (1961).

---

weak association between  $\text{K}^+$  and  $\text{IO}_3^-$ .

In conclusion, correction for ionic association of the supporting electrolyte and/or ionic reactants appears in order in the correlation between double layer structure and electrode kinetics. However, one would do well to remember the words of caution of Robinson and Stokes<sup>18</sup> when using equilibrium constants for ionic association.

---

(18) R.A. Robinson and R.H. Stokes, "Electrolyte Solutions", 2nd edition, Academic Press, Inc., New York, N.Y., 1959, pp.421-423.

---

ACKNOWLEDGMENT.— This investigation was supported by the Office of Naval Research.

TABLE I

REDUCTION OF 5 mM  $\text{KIO}_3$  IN 20 mM KOH ON MERCURY AT 25°

Added salt	E at $I = 100 \mu\text{A}\cdot\text{cm}^{-2}$ for Tafel plot V. vs. H.E. <sup>a</sup>	E - $\Delta\phi$ for corrected Tafel plot <sup>b</sup> V. vs. H.E.
none	-0.247	-0.036
0.4 M KCl	-0.105	-0.042
0.2 M $\text{K}_2\text{SO}_4$	-0.121	-0.050
0.1 M $\text{K}_4\text{Fe}(\text{CN})_6$	-0.132	-0.070

(a) H.E.  $\equiv$  hydrogen electrode in identical solution as for electrolysis but without  $\text{KIO}_3$ . (b) For  $\log I + (zF/2.3 RT)\Delta\phi = 10^3$  (I in  $\mu\text{A}\cdot\text{cm}^{-2}$ ).

Fig.1 - Corrected Tafel plots for the solutions of Table I. Correction for ionic association was made for  $K_2SO_4$  and  $K_4Fe(CN)_6$ .




FIG. 1

**TECHNICAL REPORT DISTRIBUTION LIST  
LOUISIANA STATE UNIVERSITY**

Contract Nonr 30000

NR NO. 051-258

	<u>No. Copies</u>		<u>No. Copies</u>
Commanding Officer Office of Naval Research Branch Office The John Crerar Library Building 86 East Randolph Street Chicago 1, Illinois	(1)	Air Force Office of Scientific Research (SRLT) Washington 25, D.C.	(1)
Commanding Officer Office of Naval Research Branch Office 346 Broadway New York 13, New York	(1)	Commanding Officer Diamond Ordnance Fuze Laboratories Washington 25, D.C. Attn: Technical Reference Section (ORDTL 06.33)	(1)
Commanding Officer Office of Naval Research Branch Office 1030 E. Green Street Pasadena 1, California	(1)	Office of Chief of Staff (R and D) Department of the Army Pentagon 3B516 Washington 25, D.C. Attn: Chemical Adviser	(1)
Commanding Officer Office of Naval Research Branch Office Navy #100 Fleet Post Office New York, New York	(7)	Chief, Bureau of Ships Department of the Navy Washington 25, D.C. Attn: Code 340	(2)
Director, Naval Research Laboratory Washington 25, D.C. Attn: Technical Information Officer Chemistry Division	(6) (2)	Chief, Bureau of Naval Weapons Department of the Navy Washington 25, D.C. Attn: Technical Library Code RRMA-3	(3) (1)
Chief of Naval Research Department of the Navy Washington 25, D.C. Attn: Code 425	(2)	ASTIA Document Service Center Arlington Hall Station Arlington 12, Virginia	(10)
Technical Library OASD (R and E) Pentagon Room 3E1065 Washington 25, D.C.	(1)	Director of Research Signal Corps Engineering Laboratories Fort Monmouth, New Jersey	(-)
Technical Director Research and Engineering Division Office of the Quartermaster General Department of the Army Washington 25, D.C.	(1)	Naval Radiological Defense Laboratory San Francisco 24, California Attn: Technical Library	(1)
Research Director Chemical and Plastics Division Quartermaster Research and Engineering Command Natick, Massachusetts	(1)	Naval Ordnance Test Station China Lake, California Attn: Head, Chemistry Division	(1)
		Commanding Officer Office of Ordnance Research Box CM, Duke Station Durham, North Carolina	(1)

**TECHNICAL REPORT DISTRIBUTION LIST  
LOUISIANA STATE UNIVERSITY**

Contract Nonr 30000

NR NO. 051-258

Page 2

	<u>No. Copies</u>		<u>No. Copies</u>
Brookhaven National Laboratory Chemistry Division Upton, New York	(1)	Horizons Incorporation 2891-2905, 79th Street Cleveland 4, Ohio Attn: Dr. A. J. Kolk	(1)
Atomic Energy Commission Research Division, Chemistry Branch Washington 25, D.C.	(1)	Dr. George J. Janz Department of Chemistry Rensselaer Polytechnic Institute Troy, New York	(1)
Atomic Energy Commission Library Branch Technical Information ORE Post Office Box E Oak Ridge, Tennessee	(1)	Dr. G. Barth-Wehrenalp Pennsalt Chemicals Corporation P.O. Box 4388 Philadelphia 18, Pennsylvania	(2)
U.S. Army Chemical Warfare Laboratories Technical Library Army Chemical Center, Maryland	(1)	Dr. B. R. Sundheim Department of Chemistry New York University New York 3, New York	(1)
Office of Technical Services Department of Commerce Washington 25, D.C.	(1)	Dr. H. W. Salzberg College of the City of New York Department of Chemistry New York, New York	(1)
Dr. Frank Hovorka Department of Chemistry Western Reserve University Cleveland 6, Ohio	(1)	ONR Resident Representative University of Texas P.O. Box 7786 Austin, Texas	(1)
Dr. I. M. Kolthoff Department of Chemistry University of Minnesota Minneapolis 14, Minnesota	(1)	Dr. R. N. Adams Department of Chemistry University of Kansas Lawrence, Kansas	(1)
Dr. P. Van Rysselberghe Department of Chemistry Stanford University Palo Alto, California	(1)	Dr. Morris Eisenberg Lockheed Aircraft Corporation Missiles Systems Division Sunnyvale, California	(1)
Dr. Norman Hackerman Department of Chemistry University of Texas Austin 12, Texas	(1)	Professor M. Metzger Department of Mining and Metallurgical Engineering University of Illinois Urbana, Illinois	(1)
Dr. Walter J. Hamer Electrochemistry Section National Bureau of Standards Washington 25, D.C.	(1)		

TECHNICAL REPORT DISTRIBUTION LIST  
LOUISIANA STATE UNIVERSITY

Contract Nonr 30000

NR NO. 051-258

Page 3

No. Copies

Dr. R. A. Marcus  
Department of Chemistry  
Polytechnic Institute of Brooklyn  
Brooklyn 1, New York (1)

Dr. G. J. Young  
Department of Chemistry  
Alfred University  
Alfred, New York (1)

Dr. M. S. Cohen, Chief  
Propellants Synthesis Section  
Reaction Motors Division  
Denville, New Jersey (1)

Monsanto Chemical Company  
Research and Engineering Division  
Boston 49, Massachusetts  
Attn: Mr. K. Warren Easley (1)

Commander  
Naval Ordnance Laboratory  
Corona, California  
Attn: Code 44 (1)

Dr. H. Bauer  
The University of Sydney  
Faculty of Agriculture  
Sydney, New South Wales  
Australia (1)

Dr. D.A. Brown  
Department of Chemistry  
University College  
Upper Merrion Street  
Dublin, Ireland (1)

Dr. Leonard Nanis  
School of Mines  
Columbia University  
New York 27, New York (1)